

## ОБЗОРЫ И ЛЕКЦИИ

УДК 577.833.2

## FLAVIVIRIDAE — ADDITION TO THE FAMILY

I. V. Kruglov

Federal State Budgetary Scientific Institution

«I. I. Mechnikov Scientific Research Institute of Vaccines and Serums RAMS»  
Moscow, Russian Federation

This review article analyzes and summarizes the history of the replenishment of the family Flaviviridae with new members over the last several decades on the example of the youngest genera of this family — *Hepacivirus* and *Pegivirus*.

It all started in 1966 when surgeon George Barker, who had hepatitis, had blood serum taken in containing an unknown virus. This virus was named *GBV*, by patient initials. Samples of the serum were frozen. A nucleic acid recognized as corresponding to the genomes of 2 separate virus species was isolated from the tested material in 1995. These viruses were named *GBV-A* and *GBV-B*. By this time, the hepatitis C virus had already been discovered, which was assigned to the Flaviviridae family, where a separate, third genus of *Hepacivirus* was allocated for it.

In 2010, a more distantly related virus (named *GBV-D*) was found in bats (Indian flying fox — lat. *Pteropus giganteus*). *GBV-B*, which causes acute hepatitis in experimentally infected tamarines, became the second species in the genus *Hepacivirus* to company with hepatitis C virus. The remaining GB viruses based on phylogenetic relationships, genome organization, and pathogenetic properties were proposed in 2011 to be classified as members of the fourth genus in the Flaviviridae family. This genus was named *Pegivirus* (pe — persistence, g — GB).

11 species of viruses have now been identified in the genus *Pegivirus*. They are indicated by letters in the order of the Latin alphabet — from *Pegivirus A* to *Pegivirus K*. And 14 species of viruses have now been identified in the genus *Hepacivirus*. So the story of the investigation, which began in 1966 with the discovery of the previously unknown *GBV* virus, has so far concluded with the discovery of two new genera of the family Flaviviridae. Numerous members of these two genera infect and also persist among a wide range of species belonging to different orders of the mammalian class, including *Homo Sapiens*.

**Key words:** Flaviviridae, *Hepacivirus*, *Pegivirus*, viral ecology.

В данной статье обзорного характера анализируется и обобщается история пополнения семейства Flaviviridae новыми членами за последние несколько десятилетий на примере наиболее молодых родов этого семейства — *Hepacivirus* и *Pegivirus*.

Все началось в 1966 году, когда у хирурга Джорджа Баркера, заболевшего гепатитом, была взята сыворотка крови, содержащая неизвестный вирус, который был назван *GBV*, по инициалам пациента. Образцы этой сыворотки крови были заморожены. В 1995 году из исследованного материала была выделена нуклеиновая кислота, признанная соответствующей геномам 2-х отдельных видов вирусов, получивших наименование *GBV-A* и *GBV-B*. К этому времени уже был открыт вирус гепатита С, который был отнесен в семейство Flaviviridae, где для него был выделен отдельный, третий род *Hepacivirus*.

В 2010 году более отдаленно родственный вирус (названный *GBV-D*) был обнаружен у летучих мышей (индийская летучая лисица — лат. *Pteropus giganteus*). *GBV-B*, вызывающий острый гепатит у экспериментально инфицированных тамиринов, стал вторым видом в роде *Hepacivirus*, где составил компанию вирусу гепатита С. Остальные GB вирусы на основании филогенетических взаимоотношений, организации генома и патогенетических свойств в 2011 году было предложено классифицировать как членов четвертого рода в семействе Flaviviridae, названного *Pegivirus* (pe — персистенция, g — GB).

В настоящее время идентифицировано 11 видов вирусов в составе рода *Pegivirus*. Они обозначены буквами по порядку латинского алфавита — от *Pegivirus A* до *Pegivirus K*. Идентифицировано также 14 гепацивирусов, имеющих буквенные обозначения от A до N. Так история исследования, начавшаяся в 1966 году с выявления ранее неизвестного вируса *GBV*, завершилась к настоящему времени открытием двух новых родов семейства Flaviviridae, многочисленные представители которых инфицируют, а также персистируют среди широкого спектра видов, принадлежащих к различным отрядам класса млекопитающих, включая *Homo Sapiens*.

**Ключевые слова:** Flaviviridae, *Hepacivirus*, *Pegivirus*, экология вирусов.

I. V. Kruglov

Flaviviridae — Addition to the Family

Problemy Zdorov'ya i Ekologii. 2019 Oct-Dec; Vol 62 (4): 04-10

The Flaviviridae family was separated as a separate family from the Togaviridae family in 1985, and included at that time two genera of vi-

ruses — *Flavivirus* and *Pestivirus*. After the discovery of hepatitis C virus in 1989, which has an external envelope, a linear single-stranded positive

RNA genome, based on these features as well as structural features of RNA, it was also included in the Flaviviridae family, where a separate, third genus *Hepacivirus* was established for it. In which this virus was initially in proud lone, unlike the hepatitis B virus, for example, which by that time had become a prototype virus for the whole new family of Hepadnaviridae.

But the hepatitis C virus wasn't alone for long. Yet in 1966, the surgeon George Barker, who had suffered hepatitis without HBsAgemia, came to the attention of virologists. The patient's blood, taken on the 3rd day of the icteric period, was injected to tamarins, who first developed acute hepatitis, and then — postinfection immunity, protecting them from reinfection. After that, the supposed new hepatitis virus was named *GBV*, according to the patient's initials, and the patient's serum samples were frozen for further investigation by the future virology methods [15].

This future came in 1995, when the PCR-based method, known as representative differential analysis, was used to investigate this material. As a result, a nucleic acid recognized as corresponding to the genomes of 2 separate viral species, was isolated from the tested material. These viruses were named *GBV-A* and *GBV-B* [39]. When tamarins were infected, acute hepatitis and the subsequent development of immunity caused only *GBV-B* virus. This allowed this virus to be initially seen as an etiological agent of hepatitis in this monkey species. However, the *GBV-B* virus was not detected in wildlife and its natural host is unknown [41]. Although there is a suggestion that these are the New World's wild monkeys [40]. Tamarins were resistant to *GBV-A*, and this virus was initially recognized as anthroponotic. However, a number of variants of *GBV-A* were subsequently identified in the New World's wild monkeys. The chimpanzees are also the natural host for *GBV-A*.

Also in 1995, another virus was isolated from the blood of patients with hepatitis no A — no E. The nucleotide sequence of the viral genome was quite close to the above-mentioned viruses. Therefore, this new virus has been designated as *GBV-C* [38]. *GBV-C* was especially close to *GBV-A*, which with it, as was initially thought, had 59 % identical nucleotides and 64 % identical amino acids, while with *GBV-B* it had only 30 % identical amino acids. As a result, *GBV-A* and *GBV-C* were recognized as different isolates of the *GBV-A/C* virus.

Among the many hepatitis C virus isolates, a virus was also identified, which was subsequently found to be frequent in intravenous drug abusers and people with repeated blood transfusions and in the absence of hepatitis C. Since 1995, this virus has been named the hepatitis G virus — *HGV*. However, it turned out that the nucleotide and amino acid sequences of this virus were very close

to *GBV-C* — 95 % of the nucleotide and 85 % of the amino acid sequences coincided. As a result, *HGV* and *GBV-C* were recognized as different isolates of a single virus, which at that stage received the unifying name of *HGV*.

In 2010, a more distantly related virus (named *GBV-D*) was found in bats (Indian flying fox — lat. *Pteropus giganteus*) [17]. In the result of the analysis of the nucleotide composition and genome structure, all these viruses were also classified to the same family of Flaviviridae, in which the hepatitis C virus was at that time. Initially, only *GBV-B* causing acute hepatitis in experimentally infected tamarins, was classified. It became the second species in the genus *Hepacivirus* to make up the hepatitis C virus company. The other GB viruses were not assigned to any genus in the Flaviviridae family.

Therefore, Jack Stapleton, Peter Simmonds et al. proposed in 2011 to classify *GBV-A*-like viruses, *GBV-C* and *GBV-D* on the basis of phylogenetic relationships, genome organization and pathogenetic properties as members of the fourth genus in the Flaviviridae family, named *Pegivirus* (pe — persistence, g — GB). And it was done [41].

Two species of viruses belonging to the genus *Pegivirus* were initially identified. These were *pegivirus A*, combining viruses formerly known as *GBV-A* and *GBV-C/HGV*, and *pegivirus B*, formerly known as *GBV-D*. It has been found that humans and chimpanzees are the natural hosts for the *GBV-C/HGV* subspecies of *Pegivirus A*, and chimpanzees and some New World's monkeys for the *GBV-A* subspecies of *Pegivirus A*. The first natural hosts identified for *Pegivirus B* were bats, as mentioned above.

It should be noted that, although the terminology based on the host species is not approved by the International Committee on Virus Taxonomy, *pegiviruses A* have received additional names in the classification of viruses: HPgV — for isolates obtained from humans; SPgV — for isolates obtained from New World's monkeys; SPgVcpz — for isolates obtained from chimpanzees.

Further research has identified several other species as candidates for the *Pegivirus* genus. It was a rodent pegivirus identified in white-throated woodrats (lat. *Neotoma albigula*) in 2013 [23]. Horse pegivirus was detected in these animals, respectively [22]. It was found that Theiler's disease (horse hepatitis with parenteral transmission) is caused by pegivirus too. This virus was initially named Theiler's disease virus [11]. Both horse viruses were identified in 2013. As well as the new human hepegivirus *HHPgV* or *HPgV-2*, detected in people with history of blood transfusions, in 2015 [21]. A closely related virus was identified in the graceful catshark (lat. *Proscyllium habereri*). However, the question of whether this virus be-

longs to pegiviruses or hepaciviruses has not yet been resolved [35].

At present, 11 virus species have been identified in the genus *Pegivirus* on the basis of deep sequencing technologies. These viruses differ more than 50 % in the nucleotide sequence and more than 55 % in the amino acid sequence [40]. They are indicated by letters in the order of the Latin alphabet — from *Pegivirus A* to *Pegivirus K*. For pegiviruses persisting with high frequency in the human population and initially described as GB virus C (*GBV-C*) and Hepatitis G virus (*HGV*) [26, 38], the name of the human pegivirus HPgV was subsequently proposed. Since at present there is no evidence that this virus is associated with hepatitis [41], and the surgeon GB was not infected with it. HPgV is transmitted sexually, through contact with infected blood and from mother to child. This virus has now been identified as *Pegivirus C* [40]. Another subspecies of *Pegivirus C* (*GBV-Ctro*) infects the chimpanzees [3, 9]. Initial data on the high degree of nucleotide and amino acid sequence homology of *GBV-A* and *GBV-C/HGV* viruses have not been confirmed. Thus, these viruses, initially combined into one species of *Pegivirus A*, were subsequently still divided into two species — *Pegivirus A* and *Pegivirus C*, respectively. *Pegivirus C* is not the lone human pegivirus. There is also the parenterally transmitted *HHPgV* or *HPgV-2* pegivirus [8, 21], now identified as *Pegivirus H* [40]. *Pegiviruses A, B, D, E, F, G, I, K* infect Old World's monkeys [7, 36]. At the same time, as already mentioned, *Pegivirus A* (*GBV-A*) also infects several species of New World's monkeys [10, 30], and *Pegivirus B* infects bats. *Pegivirus E* (equine pegivirus) [22] and *Pegivirus D* (Theiler's disease virus) [11] also infect horses. *Pegivirus K* also infects pigs [6]. *Pegiviruses F, G, I, J* also infect a wide range of rodent and bat species [23]. Pegiviruses, as well as hepaciviruses, differ from other flaviviruses — members of the genera *Flavivirus* and *Pestivirus*, limited ability to replicate in cell cultures. Although this barrier has been overcome for the hepatitis C virus [1, 2] as well as for the *Pegivirus C* [13].

*Pegivirus C*, formerly known as *GBV-C*, persists with high frequency in human society. But, except for the association with non-Hodgkin's lymphoma, *Pegivirus C* is not associated with the development of any identifiable disease [12, 24]. According to current information, the infection of other mammalian species with pegiviruses is persistent and nonpathogenic, except for infection of horses by *Pegivirus D* [11]. In general, this is quite typical for viruses of this genus, the name of which came from the word «persistence».

Pegiviruses have both similarities and some differences in the genome organization from vi-

ruses of the genus *Hepacivirus* and other genera of the *Flaviviridae* family (Fig. 1, 2, 3). All pegiviruses have a long 5'-untranslated region with predicted internal ribosome entry site (IRES) function. The most part of pegiviruses have a picornavirus-like IRES type I, that has no structural analogues in other flaviviruses, while some others have IRES type IV structurally similar to hepaciviruses and pestiviruses. Besides that, pegiviruses do not encode a protein similar to the nucleocapsid protein of other flaviviruses [28, 31, 41]. IRES is a structural part of the RNA, that allows to initiate cap-independent translation as part of the overall process of protein synthesis. IRES sequences were first discovered in 1988 in polyovirus (PV) [29] and encephalomyocarditis virus (EMCV) [19].

As with other flaviviruses, the pegivirus genome contains one open reading frame (ORF). The genome encodes a polyprotein that is co- and posttranslationally cuts into individual viral proteins (Fig. 1). The structural proteins common to all pegiviruses are the E1 and E2 envelope glycoproteins. The non-structural proteins NS2 — NS5B are also common to all pegiviruses. Cutting of structural proteins by cellular signal peptidases, NS2/NS3 by NS2-NS3 autoprotease and other NS proteins by NS3-NS4A protease complex is carried out in the same order as in hepaciviruses.

The size of the pegivirus genome ranges from 8900 to 11300 nucleotide bases.

Pegiviruses with a longer genome are thought to encode the additionally predicted structural proteins X and Y. Thus, although pegiviruses lack an analogue of the nucleocapsid protein characteristic for other flaviviruses, some of them probably have the predicted Y protein with an unknown function preceding E1. Several pegiviruses also probably have the predicted additional glycoprotein X, following E2 (Fig. 1).

And what is the situation with the hepaciviruses? Initially, we focused on the fact that the second species in the genus *Hepacivirus*, which accompanied the hepatitis C virus, was *GBV-B*, causing acute hepatitis in experimentally infected tamarins (see above). Then hepaciviruses were detected in bats [16], rodents [16, 23], horses [16] and dogs, and African primates [25]. Based on the results of such methods as immunoblot of the sera of infected animals, PCR and sequencing of viral genomes, J.F.Drexler et al. obtained the following initial picture of the phylogenetic relationships of potential members of the genus *Hepacivirus* with each other and with other viruses of the *Flaviviridae* family (Fig. 4) [16].

14 species of viruses have now been identified in the genus *Hepacivirus* [40]. *Hepacivirus A* is the dog's and horse hepacivirus. This virus is evolutionarily closest to the hepatitis C virus [43].

*Hepacivirus B* is a virus formerly known as *GBV-B*. *Hepacivirus C* is the hepatitis C virus. *Hepacivirus D* is the guereza hepacivirus (the eastern black-and-white colobus or the guereza (lat. *Colobus guereza*) — a species of monkey inhabiting central Africa and Ethiopia). *Hepacivirus E* is the virus formerly known as rodent hepacivirus 339. *Hepacivirus F* is the virus formerly known as rodent hepacivirus NLR07-oct70. *Hepacivirus G* corresponds to hepacivirus 1 of brown rats (lat. *Rattus norvegicus*) and *Hepacivirus H* corresponds to hepacivirus 2 of brown rats. *Hepacivirus I* corresponds to rodent hepacivirus SAR-3/RSA/2008 and *Hepacivirus J* corresponds to rodent hepacivirus RMU10-3382/GER/2010. *Hepaciviruses K, L, M* are hepaciviruses of bats formerly known as PDB-829, PDB-112 and PDB-491.1, respectively. Hepacivirus N is the cattle hepacivirus [5, 14]. And it is questionable whether hepacivirus is a new virus isolated from hoary bamboo rats (lat. *Rhizomys pruinosus*) [44]. As already mentioned above, the question of whether a virus identified in the graceful catshark (lat. *Proscyllium habereri*) belongs to *Hepaciviruses* or *Pegiviruses* has not yet been resolved [35].

*Hepaciviruses* have both similarities and some differences in the genome organization from viruses of other genera of the *Flaviviridae* family (Fig. 3). The most part of hepaciviruses contain IRES type IV in the 5'-untranslated region, however the IRES of such rodent hepaciviruses as F and J, has some homology with the corresponding sequence of pegiviruses [16, 18, 28, 40]. The hepacivirus genome sizes range from 8900 to 10500 nucleotide bases. The hepacivirus genome contains one ORF encoding a polyprotein of about 3000 amino acids (Fig. 2). The *HCV* virion contains at least three proteins: the nucleocapsid C core protein and two envelope glycoproteins, E1 and E2. An additional protein, p7 (which is supposed to have the properties of an ion channel protein important for the virus assembly), is not completely cleaved from the E2 precursor, forming E2-p7 and p7 [34]. However, it is unknown whether they are structural elements of the virion. The *GBV-B* corresponding protein, p13, is cut into p7 and p6 [42]. The genomes of other hepaciviruses have the similar organization with genomes of *HCV* and *GBV-B*, with the predicted splitting sites in the coding region, potentially producing the proteins C, E1, E2, p7/p13, NS2, NS3, NS4A, NS4B, NS5A and NS5B, comparable in the size to those of *HCV* and *GBV-B*. The only exception is the additional amino acid sequence of the NS5A guereza hepacivirus protein (*Hepacivirus D*) [25].

The cutting of viral proteins from the polyprotein in hepaciviruses occurs generally in a sim-

ilar order to pegiviruses. Structural proteins are also cut by cellular signal peptidases. The peculiarity of hepaciviruses is that they have a signal peptide peptidase involved in this process. This enzyme cuts a part of the protein already sliced from the polyprotein, unlike signal peptidases in general. Concretely, it cuts off the C-end of the nucleocapsid protein in hepaciviruses. Since pegiviruses have not the analogous nucleocapsid protein, the analogous peptide peptidase is not involved in the processing of their proteins. Cutting of non-structural proteins in hepaciviruses is the same as in pegiviruses — the NS2/NS3 sequence is cut by NS2-NS3 autoprotease and the other NS proteins are cut by the NS3-NS4A protease complex.

Hepatitis C, the infection caused by *Hepacivirus C*, is well known, and here is no need to describe it. The natural host of *Hepacivirus B*, formerly named *GBV-B*, is unknown [41], although there is a suggestion that these are wild New World's monkeys [40]. Horse hepacivirus infection is accompanied by hepatitis with a slight increase of enzyme levels [27, 30]. Quantitative PCR and histopathology of hepacivirus-infected rodents confirmed tropism to liver tissue with hepatitis C-like inflammation [16]. The pathogenicity of other hepaciviruses has not yet been sufficiently studied. However, the presence of miR-122 microRNA binding sites in the majority of hepaciviruses suggests widespread hepatotropism among species of this genus [40]. MicroRNA 122 (miR-122) is a small RNA molecule that is specifically expressed in the liver. According to the information obtained, miR-122 promotes both hepacivirus RNA replication [20, 32] and its translation [32, 33] as a result of interaction between miR-122 and the 5'-non-coding region of the viral genome [4]. This is quite typical for viruses of this genus, the name of which came from the word «hepatotropism».

However, some of the characteristics that were initially used in the classification in the long run are no strictly different for all the viruses that form these two genera. It concerns to the IRES type, the presence of miR-122 binding sites in the 5'-non-coding region of the viral genome and the virus persistence. At present, the demarcation between these genera is based on the phylogenetic difference and, above all, the presence or absence of the nucleocapsid protein.

So the story of the investigation, which began in 1966 with the discovery of the previously unknown *GBV* virus, has so far concluded with the discovery of two new genera of the family *Flaviviridae*. Numerous members of these two genera infect and also persist among a wide range of species belonging to different orders of the mammalian class, including *Homo Sapiens*.

### Pegivirus genome




Fig. 1 — Pegivirus genome organization and viral protein processing [37]

### Hepacivirus genome




Fig. 2 — Hepacivirus genome organization and viral protein processing [37]





Fig. 3 — Genome organization of typical representatives of the Flaviviridae family by the example of viruses of the genus *Flavivirus* and viral protein processing [37]





9. Birkenmeyer LG, Desai SM, Muerhoff AS, Leary TP, Simons JN, Montes CC. et al. Isolation of a GB virus-related genome from a chimpanzee. *J Med Virol.* 1998;56(1):44-51. doi: 10.1002/(sici)1096-9071(199809)56:1<44::aid-jmv8>3.0.co;2-n.
10. Bukh J, Appgar CL. Five new or recently discovered (GBV-A) virus species are indigenous to New World monkeys and may constitute a separate genus of the Flaviviridae. *Virology.* 1997;229(2):429-36. doi: 10.1006/viro.1997.8461
11. Chandriani S, Skewes-Cox P, Zhong W, Ganema DE, Diversb TJ, Van Blaricum AJ. et al. Identification of a previously undescribed divergent virus from the Flaviviridae in an outbreak of equine serum hepatitis. *Proceedings Of The National Academy Of Sciences USA.* 2013;110(15):5752-53. doi: 10.1073/pnas.1219217110
12. Chang CM, Stapleton JT, Klinzman D, McLinden JH, Purdue MP, Katki HA. et al. GBV-C infection and risk of NHL among U.S. adults. *Cancer Research.* 2014;74(19):5553-60. doi:10.1158/0008-5472.CAN-14-0209.
13. Chivero ET, Stapleton JT. Tropism of human pegivirus (formerly known as GB virus C/hepatitis G virus) and host immunomodulation: insights into a highly successful viral infection. *J Gen Virol.* 2015;96(7):1521-32. doi:10.1099/vir.0.000086.
14. Corman VM, Grundhoff A, Baechlein C, Fischer N, Gmyl A, Wolny R. et al. Highly Divergent Hepaciviruses from African Cattle. *J Virol.* 2015;89(11):5876-82. doi:10.1128/JVI.00393-15.
15. Deinhardt F, Holmes AW, Capps RB, Popper H. Studies on the transmission of human viral hepatitis to marmoset monkeys. *J Exper Med.* 1967; 125(4):673-88. doi: 10.1084/jem.125.4.673
16. Drexler JF, Corman VM, Müller MA, Lukashev AN, Gmyl A, Coutard B. et al. Evidence for Novel Hepaciviruses in Rodents. *PLoS Pathogens.* 2013;9(6): e1003438. doi:10.1371/journal.ppat.1003438.

17. Epstein JH, Quan P-L, Briese T, Street C, Jabado O, Conlan S. et al. Identification of GBV-D, a Novel GB-like Flavivirus from Old World Frugivorous Bats (*Pteropus giganteus*) in Bangladesh. *PLoS Pathogens*. 2010;6(7): e1000972. doi:10.1371/journal.ppat.1000972.
18. Honda M, Beard MR, Ping L-H, Lemon SM. A Phylogenetically Conserved Stem-Loop Structure at the 5' Border of the Internal Ribosome Entry Site of Hepatitis C Virus Is Required for Cap-Independent Viral Translation. *J Virol*. 1999;73(2):1165-74.
19. Jang SK, Kräusslich HG, Nicklin MJ, Duke GM, Palmenberg AC, Wimmer E. A segment of the 5' nontranslated region of encephalomyocarditis virus RNA directs internal entry of ribosomes during *in vitro* translation. *J Virol*. 1988;62(8):2636-43.
20. Jopling CL, Yi M, Lancaster AM, Lemon SM, Sarnow P. Modulation of hepatitis C virus RNA abundance by a liver-specific MicroRNA. *Science*. 2005;309(5740):1577-81. doi: 10.1126/science.1113329.
21. Kapoor A, Kumar A, Simmonds P, Bhuvu N, Chauhan LC, Lee B. et al. Virome Analysis of Transfusion Recipients Reveals a Novel Human Virus That Shares Genomic Features with Hepaciviruses and Pegiviruses. *Mbio*. 2015;6(5):e01466-15. doi:10.1128/mBio.01466-15.
22. Kapoor A, Simmonds P, Cullen JM, Scheel TKH, Medina JL, Giannitti F. et al. Identification of a Pegivirus (GB Virus-Like Virus) That Infects Horses. *J Virol*. 2013;87(12):7185-90. doi:10.1128/JVI.00324-13.
23. Kapoor A, Simmonds P, Scheel TKH, Hjelle B, Cullen JM, Burbelo PD. et al. Identification of Rodent Homologs of Hepatitis C Virus and Pegiviruses. *MBio*. 2013;4(2): e00216-13. doi:10.1128/mBio.00216-13.
24. Krajden M, Yu A, Braybrook H, Lai AS, Mak A, Chow R. et al. GBV-C/hepatitis G virus infection and non-Hodgkin lymphoma: a case control study. *Int J Cancer*. 2010;126(12):2885-92. doi: 10.1002/ijc.25035.
25. Lauck M, Sibley SD, Lara J, Purdy MA, Khudyakov Y, Hyeroba D. et al. A Novel Hepacivirus with an Unusually Long and Intrinsically Disordered NS5A Protein in a Wild Old World Primate. *J Virol*. 2013;87(16):8971-81. doi: 10.1128/JVI.00888-13.
26. Linnen J, Wages JJ, Zhang-Keck ZY, Fry KE, Krawczynski KZ, Alter H. et al. Molecular cloning and disease association of hepatitis G virus: a transfusion-transmissible agent. *Science*. 1996;271(5248):505-8. doi: 10.1126/science.271.5248.505.
27. Lyons S, Kapoor A, Schneider BS, Wolfe ND, Culshaw G, Brendan Corcoran B et al. Viraemic frequencies and seroprevalence of non-primate hepacivirus and equine pegiviruses in horses and other mammalian species. *J Gen Virol*. 2014;95(8):1701-11. doi: 10.1099/vir.0.065094-0.
28. Muerhoff AS, Leary TP, Simons JN, Pilot-Matias TJ, Dawson GJ, Erker JC. et al. Genomic organization of GB viruses A and B: two new members of the Flaviviridae associated with GB agent hepatitis. *J Virol*. 1995;69(9):5621-30.
29. Pelletier J, Sonenberg N. Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA. *Nature*. 1988;334(6180):320-25. doi: 10.1038/334320a0.
30. Pfaender S, Cavalleri JM, Walter S, Doerrbecker J, Campana B, Brown RJP. et al. Clinical course of infection and viral tissue tropism of hepatitis C virus-like nonprimate hepaciviruses in horses. *Hepatology*. 2015;61(2):447-59. doi: 10.1002/hep.27440.
31. Quan P-L, Firth C, Conte JM, Williamsa SH, Zambrana-Torrellob CM, Anthonya SJ. et al. Bats are a major natural reservoir for hepaciviruses and pegiviruses. *Proceedings of the National Academy of Sciences USA*. 2013;110(20):8194-99. doi: 10.1073/pnas.1303037110.
32. Roberts APE, Doidge R, Tarr AW, Jopling CL. The P body protein LSM1 contributes to stimulation of hepatitis C virus translation, but not replication, by microRNA-122. *Nucleic Acids Research*. 2014;42(2):1257-69. doi: 10.1093/nar/gkt941.
33. Roberts APE, Lewis AP. and Jopling CL. miR-122 activates hepatitis C virus translation by a specialized mechanism requiring particular RNA components. *Nucleic Acids Research*. 2011;39(17):7716-29. doi: 10.1093/nar/gkr426.
34. Shanmugam S. and Yi M. Efficiency of E2-p7 Processing Modulates Production of Infectious Hepatitis C Virus. *J Virol*. 2013;87(20):11255-66. doi: 10.1128/JVI.01807-13.
35. Shi M, Lin X-D, Vasilakis N, Tian J-H, Li C-X, Chen L-J. et al. Divergent Viruses Discovered in Arthropods and Vertebrates Revise the Evolutionary History of the Flaviviridae and Related Viruses. *J Virol*. 2016;90(2):659-69. doi: 10.1128/JVI.02036-15.
36. Sibley SD, Lauck M, Bailey AL, Hyeroba D, Tumukunde A, Weny G. et al. Discovery and Characterization of Distinct Simian Pegiviruses in Three Wild African Old World Monkey Species. *PLoS One*. 2014;9(6):e98569. doi: 10.1371/journal.pone.0098569.
37. Simmonds P, Becher P, Bukh J, Gould EA, Meyers G, Monath T. et al. ICTV Virus Taxonomy Profile: Flaviviridae. *J Gen Virol*. 2017;98(1):2-3. doi: 10.1099/jgv.0.000672.
38. Simons JN, Leary TP, Dawson GJ, Pilot-Matias TJ, Muerhoff AS, Schlauder GG. et al. Isolation of novel virus-like sequences associated with human hepatitis. *Nature Medicine*. 1995;1(6):564-69. doi: 10.1038/nm0695-564.
39. Simons JN, Pilot-Matias TJ, Leary TP, Dawson GJ, Desai SM, Schlauder GG. et al. Identification of two flavivirus-like genomes in the GB hepatitis agent. *Proceedings of the National Academy of Sciences USA*. 1995;92(8):3401-3405. doi: 10.1073/pnas.92.8.3401.
40. Smith DB, Becher P, Bukh J, Gould EA, Meyers G, Monath T. et al. Proposed update to the taxonomy of the genera Hepacivirus and Pegivirus within the Flaviviridae family. *J Gen Virol*. 2016;97(11):2894-2907. doi: 10.1099/jgv.0.000612.
41. Stapleton JT, Fong S, Muerhoff AS, Bukh J, Simmonds P. The GB viruses: a review and proposed classification of GBV-A, GBV-C (HGV), and GBV-D in genus Pegivirus within the family Flaviviridae. *J Gen Virol*. 2011;92(2):233-46. doi: 10.1099/vir.0.027490-0.
42. Takikawa S, Engle RE, Emerson SU, Purcell RH, Claire MS, Bukh J. Functional analyses of GB virus B p13 protein: Development of a recombinant GB virus B hepatitis virus with a p7 protein. *Proceedings of the National Academy of Sciences USA*. 2006;103(9):3345-50. doi: 10.1073/pnas.0511297103.
43. Theze J, Lowes S, Parker J, Pybus OG. Evolutionary and Phylogenetic Analysis of the Hepaciviruses and Pegiviruses. *Genome Biol Evol*. 2015;7(11):2996-3008. doi:10.1093/gbe/evv202.
44. Van Nguyen D, Van Nguyen C, Bonsall D, Tri Ngo T, Carrique-Mas J, Hong Pham A. et al. Detection and Characterization of Homologues of Human Hepatitis Viruses and Pegiviruses in Rodents and Bats in Vietnam. *Viruses*. 2018;10(3):102. doi:10.3390/v10030102.

#### Адрес для корреспонденции

115088, Российская Федерация,  
Москва, ул. Первая Дубровская, 15,  
ФГБНУ «Научно-исследовательский институт вакцин и сывороток им. И.И.Мечникова»,  
Отдел вирусологии им. О.О.Анжaparидзе,  
Лаборатория молекулярно-биологической технологии,  
Тел. моб.: +7 917 5329418,  
e-mail: [kruglov15@inbox.ru](mailto:kruglov15@inbox.ru)  
Круглов Игорь Вячеславович

#### Сведения об авторах

Круглов И.В., д.б.н., ведущий научный сотрудник, лаборатория молекулярно-биологической технологии ФГБНУ «Научно-исследовательский институт вакцин и сывороток им. И.И.Мечникова».

#### Address for correspondence

115088, Russian Federation,  
Moscow, Pervaya Dubrovskaya Str., 15,  
Mechnikov Research Institute of Vaccines & Sera,  
O.G.Andzhaparidze Department of Virology,  
Laboratory of Molecular Biological Technology,  
Mob. tel.: +7 917 5329418,  
e-mail: [kruglov15@inbox.ru](mailto:kruglov15@inbox.ru)  
Kruglov Igor Vyacheslavovich

#### Information about the authors

Kruglov I.V., Dr. Sci., Leader Researcher, Laboratory of Molecular Biological Technology, Mechnikov Research Institute of Vaccines & Sera.

Поступила 06.11.2019